

The Run II Physics Program

John Womersley

Fermi National Accelerator Laboratory, Batavia, Illinois

Representing the CDF and DØ collaborations



The Run II physics program

- A broad physics program which evolves as a function of luminosity
 - There is interesting physics at all luminosities, starting now with 50-100 pb⁻¹ and continuing through 0.3, 1, 2, 5, 10, 15 fb⁻¹
- This physics program has begun
- The goal of the collaborations is to
 - Maximize this physics program
 - Exploit the full potential of the world's highest energy collider and the large investments we have made in the accelerator and detectors
 - Lay a firm foundation for the LHC and for future initiatives at the TeV scale
 - Attract and train the best students in the field
 - Clarify physics requirements
 - An international program in the US - groundwork for the future





Big Questions at the Electroweak Scale

- The Tevatron is the only accelerator in operation that can help to answer
 - What is the structure and what are the symmetries of space-time?
 - Why is the weak force weak?
 - What is cosmic dark matter made of?



About six to seven times more mass in the universe ($27 \pm 4\%$) than there is baryonic matter ($4.4 \pm 0.4\%$)



What is this stuff? How can we get a firmer understanding of it?

Accelerators

- Run II is the only opportunity to make such a major discovery at an accelerator in the United States



The program

- **The Run II Physics program**
 - **Confront the standard model through precise measurements**
 - **The strong interaction, the quark mixing matrix, the electroweak force and the top quark**
 - **Directly search for particles and forces not yet known**
 - **Both those predicted (Higgs, supersymmetry, dark matter, extra dimensions) and those that would come as a surprise**
- **The program was developed in a series of workshops between 1998 and 2000**
 - <http://fnth37.fnal.gov/run2.html>
- **The program stretches from the GeV scale to the TeV scale**
- **Here I can attempt only a superficial survey and will concentrate on the physics that gains most from luminosity**
 - **Illustration of the breadth of the program:
110 talks from CDF and DØ at APS/DPF meeting in April**



Two Worldwide Collaborations

More than 50% non-US: a central part of the world HEP program



**12 countries, 59 institutions
706 physicists**

John Womersley, slide 5




**18 countries, 78 institutions
664 physicists**

19 ~~80~~ 670




Operations Status

- Both experiments are operating well and recording physics quality data with high ($\sim 90\%$) efficiency
- 100-140 pb^{-1} being used for analysis for summer 2003
- Data are being reconstructed within a few days






The Top Quark

- Why, alone among the elementary fermions, does the top quark couple strongly to the Higgs field?
 - Is nature giving us a hint here?
 - Is the mechanism of fermion mass generation indeed the same as that of EW symmetry breaking?
 - The top is a window to the origin of fermion masses
- The Tevatron Collider is the world's only source of top quarks
- We are measuring its
 - Mass
 - Production cross section
 - Spin
 - Through top-antitop spin correlations
 - Electroweak properties
 - Through single top production
- Any surprises, anomalies?




The Run II Top Physics Program has begun






The top quark rediscovered, 2003



Cross section


CDF dileptons $\sigma = 13.2 \pm 5.9_{stat} \pm 1.5_{sys} \text{ pb}$
CDF $l + jets$ $\sigma = 5.3 \pm 1.9_{stat} \pm 0.8_{sys} \pm 0.8_{lum} \text{ pb}$
DØ (comb.) $\sigma = 8.4^{+4.5}_{-3.7} (stat)^{+5.3}_{-3.5} (syst) \pm 0.8(lumi) \text{ pb}$

DØ Preliminary



CDF II Preliminary (72 pb⁻¹)

$M_{TOP} = 171.2 \pm 13.4 \pm 9.9 \text{ GeV}/c^2$




CDF mass

$M_{top} = 171.2^{+14.4}_{-12.5 \text{ stat}} \pm 9.9_{sys} \text{ GeV}/c^2$



Top mass

- We can look forward to improved precision on m_t in the near future
 - More data (few hundred pb^{-1})
 - Expect ~ 500 b-tagged lepton+jets events per experiment per fb^{-1}
 - cf. World total at end of Run I ~ 50
- Improved techniques
 - e.g. new DØ Run I mass measurement is equivalent to a factor 2.4 increase in statistics:



$$m_{top} = 179.9 \pm 3.6 \text{ (stat) GeV}/c^2 \text{ (5.6 GeV from PRD 58 052001, 1998)}$$

Improved top mass measurements help to constrain the Higgs mass

| Δm_t | $l + \text{jets}$ | dilepton |
|----------------------|-----------------------|-----------------------|
| 2 fb^{-1} | $\pm 2.7 \text{ GeV}$ | $\pm 2.8 \text{ GeV}$ |
| 10 fb^{-1} | $\pm 1.6 \text{ GeV}$ | $\pm 1.6 \text{ GeV}$ |

per experiment, using the
"classic" technique

[from M. Grunewald et al.,
hep-ph/0111217 (2001)]

$$\Delta m_H / \Delta m_t \sim 50 \text{ GeV} / 4 \text{ GeV}$$



Top physics program

- Precise knowledge of m_t (~ 1 GeV) will be very useful even after a light Higgs is discovered

- Is it H_{SM} or SUSY h ?
- Constrain the stop sector:
[M. Beneke et al., hep-ph/0003033]

- Single top production

- The way to measure top width
- So far unobserved
- With $\sim 1 \text{ fb}^{-1}$ should be able to see signals for both s and t-channel production (have different sensitivity to new physics)

| | $\Delta\sigma$ (s) | $\Delta V_{tb} $ (s) | $\Delta\sigma$ (t) | $\Delta V_{tb} $ (t) |
|---------------------------|--------------------|----------------------|--------------------|----------------------|
| 2 fb⁻¹ | 21% | 12% | 12% | 10% |
| 10 fb⁻¹ | 9% | 6% | 5% | 8% |


[scaled from T. Stelzer, Z. Sullivan and S. Willenbrock, Phys. Rev. **D58**, 094021 (1998)]

- Top-antitop spin correlations

- With 2fb^{-1} , distinguish spin-1/2 from spin-0 but only at the 2σ level




- New physics

- $t\bar{t}$ mass, top p_T , rare decays and nonstandard decays, anomalous single top ...



Electroweak Physics

- In Run II we will complement direct searches for new phenomena with indirect probes
 - New particles and forces can be seen indirectly through their effects on electroweak observables.
 - Tightest constraints come from improved determination of the masses of the W and the top quark.
- Both experiments have preliminary results from Run II samples of W and Z candidates:



Prospects for W mass

Current knowledge of m_W


- **hadron colliders:**
 - **$80\,454 \pm 59$ MeV**
- **World (dominated by LEP)**
 - **$80\,451 \pm 33$ MeV**

Run II prospects

(per experiment)

| | Δm_W |
|---------------------------------------|--------------------------------|
| 2 fb^{-1} | ± 27 MeV |
| 10 fb^{-1} | ± 18 MeV |

[from M. Grunewald et al.,
hep-ph/0111217 (2001)]




We have shown we can measure the W mass precisely at the Tevatron, but to improve on LEP will require $\sim \text{fb}^{-1}$ datasets - not a short term goal

$$dm_H/dm_W \sim 50 \text{ GeV}/25 \text{ MeV}$$




Other electroweak measurements

- Forward-backward asymmetry A_{FB} in $Z \rightarrow ee$



CDF – Paper in preparation



Projection for 10 fb^{-1}

- measure effective $\sin^2\theta_w$ to 0.0002 (10fb^{-1}) and test γ^*/Z interference at \sqrt{s} much greater than LEP
- Other electroweak measurements
 - Multiboson production (test gauge couplings)
 - Boson plus jets
 - ...






QCD


- No one doubts that QCD describes the strong interaction between quarks and gluons
 - Its effects are all around us:
 - masses of hadrons (stars and planets)
 - But it is not an easy theory to work with
- Use the Tevatron to
 - Test QCD itself
 - Understand some outstanding puzzles from Run I
 - Develop the expertise to calculate, and confidence in, the backgrounds to new physics
 - Excellent interaction between the experimental and phenomenology communities



Some QCD Physics goals for Run II





Run I jet data already used in CTEQ6 and MRST2001 parton distribution fits; complements HERA's kinematic range





b-jet cross section
Important background to new physics

Jets in Run II



DØ Run II Preliminary



Searches for New Physics

- The Tevatron, as the world's highest energy collider, is the most likely place to directly discover a new particle or force
- We know the SM is incomplete
 - Most popular extension: supersymmetry
- Predicts multiple Higgs bosons, strongly interacting squarks and gluinos, and electroweakly interacting sleptons, charginos and neutralinos
 - masses depend on unknown parameters, expected to be 100 GeV - 1 TeV

**Lightest neutralino is a good candidate for cosmic dark matter
Potentially discoverable at the Tevatron**




Supersymmetry signatures

- Squarks and gluinos are the most copiously produced SUSY particles
- As long as R-parity is conserved, cannot decay to normal particles
 - Jets plus missing transverse energy signatures


Make dark matter at the Tevatron!

Detect its escape from the detector

Possible decay chains always end in the LSP



Missing E_T backgrounds




Searching for squarks and gluinos

Run I


$$M_{\tilde{g}} > 300 \text{ GeV}/c^2 \quad M_{\tilde{q}} \approx M_{\tilde{g}}$$

$$M_{\tilde{g}} > 195 \text{ GeV}/c^2$$

**With 2 fb⁻¹:
Reach in gluino
mass ~ 400 GeV**




Run II analysis has begun:





**High mE_T
candidate
event**

DØ



Chargino/neutralino production

- “Golden” signature
 - Three leptons
 - very low standard model backgrounds
- This channel becomes increasingly important as squark/gluino production reaches its kinematic limits (masses ~ 500 GeV)
- Reach on χ^\pm mass, $2\text{fb}^{-1} \sim 180$ GeV ($\tan \beta = 2, \mu < 0$)
 ~ 150 GeV (large $\tan \beta$)





Searches have begun.
 So far number of events is consistent with expectations — we need a lot more data, but the tools are in place

Run II Trilepton candidate



Other Searches at the Tevatron

- Other Tevatron search channels for SUSY
 - GMSB \rightarrow Missing E_T + photon(s)
 - Stop, sbottom
 - RPV signatures
- Searches for other new phenomena
 - leptoquarks, dijet resonances, W', Z' , massive stable particles, doubly charged particles...





Several search results already comparable or better than Run I

CDF Run II $Z' > 650 \text{ GeV}/c^2$



Extra Dimensions

- Run II is also testing the new and exciting idea of extra dimensions
 - Can gravity propagate in more than four dimensions of space-time?
 - If these dimensions are not open to the other SM particles and forces, we would not perceive them
 - Particle physics experiments at the TeV scale could see effects (direct and indirect)
 - Measure the structure of space-time!




| | GRW | HLZ for n: | | Hewett |
|------|------|------------|------|----------------|
| | | 2 | 7 | $\lambda = +1$ |
| diEM | 1.12 | 1.16 | 0.89 | 1.00 |
| diMU | 0.79 | 0.68 | 0.63 | 0.71 |

DØ
Run II
Preliminary

With 300 pb^{-1} , we probe up to 1.6 TeV
With 2 fb^{-1} , we probe up to 2 TeV








*There are more things in
heaven and earth, Horatio,
Than are dreamt of in your
philosophy.*

Signature-based searches

We need to ensure that our searches are not constrained by our preconceptions of what might be “out there.”



CDF dilepton top events



Follow up anomalies in Run I data, and set model-independent limits

“Sleuth” framework used very successfully by DØ

Phys. Rev. D {62} 92004 (2000)

John Womersley, slide 24




The Higgs Boson

- In the Standard model, the weak force is weak because the W and Z gain mass from a scalar field that fills the universe
- The same field is responsible for the mass of the fundamental fermions
- If it exists, we can excite the field and observe its quanta in the lab
 - **The Higgs boson**
 - Last piece of the SM
 - Key to understanding beyond-the-SM physics like supersymmetry: a light Higgs is a basic prediction of SUSY
- All the properties of the Higgs are fixed in the SM with the exception of its own mass: simulations have no free parameters




Higgs Hunting at the Tevatron

- For any given Higgs mass, the production cross section and decays are all calculable within the Standard Model
- Inclusive Higgs cross section is quite high: $\sim 1\text{pb}$
 - for masses below $\sim 140\text{ GeV}$, the dominant decay is $H \rightarrow b\bar{b}$ which is swamped by background
 - at higher masses, can use inclusive production plus WW decays
- The best bet below $\sim 140\text{ GeV}$ appears to be associated production of H plus a W or Z
 - leptonic decays of W/Z help give the needed background rejection
 - cross section $\sim 0.2\text{ pb}$



Preliminary update on Higgs sensitivity

- A CDF + DØ joint study has updated the “famous” reach plot:




- Concentrate on low mass region (favored both in SM and SUSY)
- The Higgs reach at the Tevatron appears to be at least as good as was projected four years ago
- Our understanding will continue to improve




SUSY Higgs Production at the Tevatron

- $bb(h/H/A)$ enhanced at large $\tan \beta$:




- $\sigma \sim 1 \text{ pb}$ for $\tan \beta = 30$ and $m_h = 130 \text{ GeV}$

**CDF Run I analysis (4 jets, 3 b tags)
sensitive to $\tan \beta > 60$**



$bb(h/A) \rightarrow 4b$



**5 fb^{-1}
 $m_A = 140 \text{ GeV}$,
 $\tan \beta = 30$**




What if we see nothing?


Exclusion of a Higgs would itself be a very important result for the Tevatron

- In the SM, can exclude most of the allowed mass range with 10 fb^{-1}
- In the MSSM, can potentially exclude all the remaining parameter space with 5 - 10 fb^{-1}
- Would certainly make life “interesting” for SUSY at the TeV scale


Note: these plots do not yet take account of the Higgs reach update



95% exclusion



5σ discovery



Exclusion and discovery for SUSY Higgs sector, maximal stop mixing, sparticle masses = 1 TeV

Conclusions

- The Run II physics program has begun
 - We now have more data than Run I, taken at a higher CM energy with better detectors
- The combination of highest accelerator energy, excellent detectors, enthusiastic collaborations, and data samples that double every year guarantees interesting and important new physics results at every step.
- Each step answers important questions, and each step leads on to the next
- The collaborations are committed to
 - Maximize this physics program
 - Exploit the full potential of the world's highest energy collider and the large investments we have made in the accelerator and detectors
 - Lay a firm foundation for the LHC and for future initiatives at the TeV scale



- **We believe these goals**
 - **call for a robust effort to deliver luminosity of order 10 fb^{-1} by the time LHC starts to produce physics**
 - **require the silicon detectors be upgraded to exploit the full luminosity**

